

Bibliography – In-Check Dial and In-Check Kit

1. Amirav I, Newhouse M.T. Measurement of PIF with the in-check dial device to simulate low resistance (-Diskus) and high resistance (-Turbohaler) DPIs in children with asthma. *European Respiratory Journal*, Vol 16, Suppl. 31, August 2000, P3811 (540s).
2. Assi K, Chrystyn H. The different resistance of dry powder inhalers (DPIs), Presented at the American Thoracic Society International Conference, San Francisco, 2001, in *Am. J. Resp. Crit. Care Med.*, 163(5), B52, Poster K124, 2001.
3. Bell J.R.N, Spencer D.W. Variations in Inspiratory flow rate achieved through Dry Powder Inhalers when minimal explanation is offered to the patient. *Aerosol Society Meeting, Drug Delivery to the Lungs XI*, 11 & 12th December 2000. ISBN 0 9529777 5 3.
4. Bell J.R, Spencer D.W, Dickens C. In-Check® Dial®: Accurate Reproduction of Internal Resistance facilitates objective analysis of patient inhaler technique. *The 3rd Triennial World Asthma Meeting, Chicago, USA, July 2001*.
5. Bell J.R, Spencer D.W. Peak Inspiratory flow achieved through dry powder inhalers when minimal explanation is offered to the user. *The 3rd Triennial World Asthma Meeting, Chicago, USA, July 2001*.
6. Brajnik V.A, Sakharova G.M, Makretskaja O.N, Belevsky A S, Chuchalin A.G. The role of Inspiratory flow measurement in choosing suitable inhaler device in asthmatic patients. *European Respiratory Journal*, Vol 16, Suppl. 31, August 2000, P2206 (316s)
7. Broeders M., Molema J., Folgering. H. Peak inspiratory flow rate and the slope of the inhalation profile in dry powder inhalers (DPI). *European Respiratory Journal*, Vol 18, Suppl. 33, September 2001, P532 (69S)
8. Chrystyn H, Corrado OJ, Brownlee K, Kanthapillai P, Pearson s, Chetcuti P, Tarsin W. Inhalation rates of asthmatic children (CHILD) and adults and of chronic obstructive pulmonary disease (COPD) patients through inhalers. *American Journal of Respiratory and Critical Care Medicine*. Vol 165 (8), pA190, 2002.
9. Chrystyn H. Is inhalation rate important for a dry powder inhaler ? using the In-Check Dial to identify these rates *Respiratory Medicine* Vol 97 (2003) 2 : 181-187.
10. Conlin T.P. Inspiratory flow rates of school children who use metered dose inhalers (abstract). *Respir Care* 2002;47(9):1074.
11. Conlin T. Inspiratory flow rates of children who use metered dose inhalers. Poster P-21, *National Conference on Asthma 2003, Washington, D.C.*
12. Conlin T. Assess usefulness of In-Check DIAL (Ferraris Medical) in children who use metered dose inhalers Poster P-82, *National Conference on Asthma 2003, Washington, D.C.*
13. Da Silva SL, Michelena T, Pregal AL et al Asthma therapeutics: selecting the inhaling device *European Journal of Allergy and Clinical Immunology* 2002 Vol 57 Supplement 73 A4

14. Davis K, Wilson J, De Soyza A, Higgins B. Inspiratory flow in COPD patients; implications for inhaler prescribing patterns *Thorax* Vol 56, Supplement 3, Dec 2001, P147
15. Dempsey OJ, Fowler, SJ, Wilson A, Kennedy G, Lipworth BJ. Effects of Adding Either a Leukotriene Receptor Antagonist or Low-Dose Theophylline to a Low or Medium Dose of Inhaled Corticosteroid in Patients With Persistent Asthma *Chest* 2002; 122:151–159
16. De Ruiter M.J, Eysvogel M.M.M, Van der Palen J. Peak Inspiratory flow through Diskus® and Turbuhaler® in COPD. American Thoracic Society, San Francisco 2001, Poster Presentations.
17. Desager K.N., Geldhof J., Claes R., De Backer W. Measurement of inspiratory flow through 3 different dry powder inhalation devices using In-check in asthmatic children. *European Respiratory Journal*, Vol 18, Suppl. 33, September 2001, P3291 (493S)
18. Emeryk A, Bartkowiak-Emeryk M, Kokot M, Czerwinska-Pawluk I. Assessment of Inspiratory flow rate (PIFR) is advisable before choosing dry powder inhaler for treatment of asthmatic children. *European Respiratory Journal*, Vol 16, Suppl. 31, August 2000, P3807 (540s)
19. Emeryk A, Bartkowiak-Emeryk M, Czerwinska-Pawluk I., Postepski J. Usefulness of peak inspiratory flow measurement for selection of dry powder inhalers in asthmatic children. [Article in Polish] *Pol Merkuriusz Lek* 2000 Oct;9(52):667-70
20. Emeryk A., Emeryk M., Kokot M. Selection of the dry powder inhaler based on peak inspiratory measurements in the pre-selected group of patients with bronchial obstruction. *European Respiratory Journal*, Vol 18, Suppl. 33, September 2001, P703 (99S)
21. Essig K.M, Spohn S, Petro W. Inspiratory flows measured by 'In-Check.' *American Journal of Respiratory and Critical Care Medicine*, Vol 161, No 3, March 2000, A:690.
22. Florkiewicz E. Not the mean value of peak inspiratory flow (PIF), but the number of patients who are able to gain minimal and/or optimal peak inspiratory flow with resistance represented dry powder inhaler, is the proper measure of its utility. *European Respiratory Journal*, Vol 18, Suppl. 33, September 2001, P1655 (241S)
23. Fonseca J.A., Winck J.C., Eusebio E., Viana P., Almeida J., Marques J.A. Peak inspiratory flow in COPD: implications for the use of dry powder inhalers. *European Respiratory Journal*, Vol 18, Suppl. 33, September 2001, P901 (134S)
24. Garcia-Marcos Alvarez L., Martinez Torres A, Huillen Perez JJ., Martinez Victoria A. Peak inspiratory flow using two different inhalers and a new portable device. *An. Esp. Pediatr.* 2001 Feb;54(2):110-113
25. Giner J, Calpena M, Casan P, Sanchis J. Maximum Inspiratory flow (MIF) in patients using dry powder devices (DPD) for inhalation therapy. American Thoracic Society, San Francisco 2001, Poster presentations.
26. Han M Y Jung K H Estimation of the possible age for using various types of inhaler *European Journal of Allergy and Clinical Immunology* 2002 Vol 57 Supplement 73 A618

27. Huchon G, Dessanges J.F, Pin I, Aiache J.-M, Barthe J, Perilliat J.-G. Mesure du debit inspiratoire de pointe sur une population de 7 135 patients présentant un asthme ou une BPCO lors de la prescription d'un dispositif pour inhalation. 5 Congrès de Pneumologie de Langue Française, Article 34.
28. Huchon G.J., Dessanges J.F., Pin I., Aiache J.-M., Barthe J., Perilliat J.G. Peak inspiratory flow (PIF) when using a dry powder inhaler in asthmatics and COPD. *European Respiratory Journal*, Vol 18, Suppl. 33, September 2001, P702 (99S)
29. Kamps A.W.A, Brand P.L.P, Vermue N.A, Roorda R.J. Home monitoring of Peak Inspiratory flow through dry powder inhalers in children with asthma. *American Thoracic Society*, San Francisco 2001, Poster presentations.
30. Kokot M. Selection of a personal dry powder inhaler based on peak inspiratory flow measurements in selected groups of patients with bronchial obstruction. [Article in Polish] *Pol Merkuriusz Lek* 2000 Oct;9(52):672-676
31. Makieda D, Ishida S, Orii T, Okano Y The Influence Of Peak Inspiratory Flow And Inhalation Technique On Inhaled Steroid To Patients With Bronchial Asthma *Yakugaku Zasshi*, 123(3),211-215 (2003)
32. Nsour W.M., Alldred A., Corrado O.J., Chrystyn H. Measurement of peak inhalation rates with an In-Check Meter to identify an elderly patient's ability to use a Turbuhaler. *Respiratory Medicine* Vol 95, 2001 965-968
33. Peacock K, Diggory P, Jones V, Trembl J. Ability of elderly patients to generate adequate inspiratory flow rates to use Turbuhaler, Clickhaler, Autohaler or Easibreathe Inhalers. *Age and Ageing* 2000, 29:s2.
34. Pearce L. Do Health Professionals Have Sufficient Knowledge And Skill To Teach Optimum Inspiratory Flow (OIF)? A Study Using The In-Check Dial™ (ICD) To Evaluate Inspiratory Technique *American Journal of Respiratory and Critical Care Medicine*. 2002;165 (8):B3
35. Pearce L. Revisiting Inhalers – Selecting an Inhaler *Practice Nursing* 2002, Volume 13, No 12, p561
36. Spencer DW, Dickens C, Bell J.R.N. Improving drug delivery from common inhalers: Accurate production of internal resistance facilitates practical analysis of patient technique. *Aerosol Society Meeting, Drug Delivery to the Lungs XI*, 11 & 12th December 2000. ISBN 0 9529777 5 3.
37. Stenhouwer-de Gooijer M, Artes H G M, van der Ent C K Het gebruik van de PIF-Meter bij de inhalatorkeuze (The Use of the PIF Meter in Choosing an Inhaler) *Longkruid* 2001;182:6-9
38. Tarsin W., Assi K., Corrado O.J, Chrystyn H. Measurement of patient inhalation rates for different dry powder inhalers (DPIs) using the In-Check DIAL *Thorax* Vol 55, Supplement 3, Dec 2000, P154

39. Tarsin W, Brownlee K, Chetcuti P, Corrado O.J, Chrystyn H. Use of the In-Check device to identify inhalation rates through dry powder inhalers (DPIs). American Thoracic Society, San Francisco 2001, Poster presentations.
40. Tarsin W., Corrado O.J., Brownlee K., Kanthapillai P., Pearson S., Chetcuti P., Chrystyn H. Inter-individual variability of peak inhalation rate through different inhalers by asthmatic children and adults and those with COPD. European Respiratory Journal, Vol 18, Suppl. 33, September 2001, P900 (133S)
41. Tarsin W., Hawksworth G., Haslam G. H., Chrystyn H. Inter- and Intra- variability of inhalation rates when patients use their dry powder inhalers (DPIs) Thorax Vol 56, Supplement 3, Dec 2001, P60
42. Tarsin WY, Al-Showair RAM, Corrado OJ, Pearson SB, Chrystyn H. Peak inhalation rates (PIFRs) of COPD patients when using a Metered Dose Inhaler (MDI) and recently introduced Dry Powder Inhalers (DPI). Thorax 2003 ; 58 (S3) iii73
43. Van den Brande P, Verniest T, Verwerft J, Liesenborghs A, Vancayzeele, Decramer M. Inspiratory Flows in Elderly COPD Patients Chest 2002 122: 4 137S
44. Van der Palen J. Use of Peak Inspiratory flow for Inhaler Selection. European Respiratory Journal, Vol 14, Suppl. 30, October 1999 P2282 (339s)
45. Van der Palen J. Peak inspiratory flow through Diskus and Turbuhaler, measured by means of a peak inspiratory flow meter (In-Check DIAL[®]) Respiratory Medicine Vol 97 (2003) 3: 285-289